

Entomological Science (2011) 14 (3): 366-369

SHORT COMMUNICATION

Incidence of *Anagrus obscurus* (Hymenoptera: Mymaridae) egg parasitism on *Calopteryx haemorrhoidalis* and *Platycnemis pennipes* (Odonata: Calopterygidae: Platycnemididae) in Italy

Serena SANTOLAMAZZA¹, Enrique BAQUERO³ and Adolfo CORDERO-RIVERA²

¹Misión Biológica de Galicia (CSIC), Departamento de Genética Vegetal, Pontevedra, ²Grupo ECOEVO, Universidade de Vigo, EUET Forestal, Campus Universitario, Pontevedra, and ³Universidad de Navarra, Facultad de Ciencias, Departamento de Zoología y Ecología, Pamplona, Spain

Abstract

Very little is known about the incidence of egg parasitoids in odonates, perhaps because Odonata eggs are well protected by stems or leaves, sometimes below water. In Central Italy (Pontecorvo, Frosinone Province) two damselflies, *Calopteryx haemorrhoidalis* and *Platycnemis pennipes*, occur in high densities. In August 2007 we collected 30 stems of the aquatic plant *Potamogeton* sp. used as substrate for oviposition and incubated eggs in the laboratory. Most stems (24 for *C. haemorrhoidalis* and 23 for *P. pennipes*) contained Odonata eggs. Parasitoids emerged from 12 stems, with a mean parasitism of 2% for *C. haemorrhoidalis* and 6% for *P. pennipes*, and a maximum of 14% and 50%, respectively. Furthermore, we observed egg-laying of 19 females of *C. haemorrhoidalis* and 11 of *P. pennipes*, and marked the stems where oviposition was observed. Clutches remained in the river for five days and were then collected and incubated. Parasitoids emerged from 11 of 30 stems, with an average parasitism of 8% for *C. haemorrhoidalis* and 3% for *P. pennipes* (maximums of 50% and 29%, respectively). All parasitoids belonged to the family Mymaridae, and were identified as *Anagrus (Anagrus) obscurus* Förster, 1861, *sensu* Soyka, 1955. This is the first time that this species is described as an egg parasitoid of odonates, and that the egg parasitoid of *C. haemorrhoidalis* and egg parasitoid of odonates.

Key words: damselflies, egg parasitoid, endophytic oviposition, parasitism rate.

It is widely known that most insect species are hosts of one or more parasitoids, either in the egg, larval, pupal or adult stage (Askew 1971). Egg parasitoids are hymenopterous species specialized in searching, locating and parasitizing recently laid eggs (Godfray 1994). They are idiobiont organisms (i.e. those prevent further development of the host after initial parasitization) whose larvae develop inside the host egg, consuming the yolk, which greatly limits their adult size. Given that the immune system of insect embryos is still not developed, eggs are especially vulnerable to parasitoid attack (Beckage 1985). For this reason, insect eggs are commonly protected from parasitoids by oothecae, by protecting coverings such as scales, setae, feces, silk or spumaline, by increasing the hardness and thickness of egg chorion, or by introducing eggs into plant tissues (Gross 1993).

Literature on Odonata egg parasitism is scarce and data on parasitism rate is just occasionally presented, perhaps because odonate eggs are commonly laid in stems, leaves or other rigid structures, sometimes below water, making them well protected against egg parasitoids. Predation of endophytic odonate eggs by other animals is rarely reported, with the exception of predation by nymphs of water mites or the eggs observed in the stomach of insectivorous fishes (Corbet 1999). The only detailed information on egg mortality of endophytic dragonflies is associated with parasitism by micro-hymenopterans belonging to the superfamily

Correspondence: Serena Santolamazza, Misión Biológica de Galicia (CSIC), Departamento de Genética Vegetal, P.O. Box 28, 36080 Pontevedra, Spain. Email: anaphes@gmail.com Received 14 December 2010; accepted 18 January 2011.



Figure 1 Underwater oviposition by *Calopteryx haemorrhoidalis* on *Potamogeton* sp. plants.

Chalcidoidea, in which four families (Eulophidae, Mymaridae, Scelionidae, Trichogrammatidae) and ten genera are represented (Corbet 1999). In the current study, we identified the parasitoid of the eggs of *Calopteryx haemorrhoidalis* Vander Linden (Odonata) and *Platycnemis pennipes* Pallas (Odonata), and evaluated the parasitism rate achieved.

The study was conducted in August 2007 in central Italy, 120 km from Rome (Pontecorvo, Frosinone Province), in the stream Forma Quesa (41°25'35"N, 13°39'11"E), where both C. haemorrhoidalis and P. pennipes have large populations. The stream had a rocky bottom, with muddy and sandy areas, with a well-developed riparian forest dominated by Populus alba (Salicaceae), Alnus glutinosa (Betulaceae), Acer monspessulanus (Sapindaceae) and Salix sp. (Salicaceae). In order to study egg parasitism of C. haemor*rhoidalis* and *P. pennipes*, we collected 30 stems of the aquatic plant Potamogeton sp. (Potamogetonaceae), the main egg-laying substrate in the studied locality, with signs of having been used for oviposition by odonates. Such stems were floating or up to 5 cm below the water surface. While some C. haemorrhoidalis females submerge to lay eggs (Fig. 1), P. pennipes females only introduce the abdomen under the water. The collected stems were maintained in plastic boxes with river water and kept at room temperature (approx. 26°C) until either the parasitoid emerged or the larvae of the odonates hatched. Because eggs of Calopteryx and Platycnemis greatly differ in length (1.3 and 0.9 mm, respectively), emerged parasitoids were easily assigned to each odonate depending on their size. Unhatched eggs were opened under a dissecting microscope. Emerged parasitoids were fixed in 90% ethanol for taxonomic identification. Parasitism rate was calculated as the ratio between parasitized eggs and the total amount of eggs of each odonate.

Furthermore, to investigate the window of vulnerability of fresh eggs, we observed egg-laying of 19 females of *C. haemorrhoidalis* and 11 of *P. pennipes*, and marked the stems were oviposition was observed. Clutches remained in the river for five days and were then collected and incubated. We chose this time interval because older eggs would be unsuitable for parasitism, considering that a significant negative correlation exists between the duration of the host development (approximately two weeks for both odonate embryos) and the percentage of host eggs successfully parasitized (Strand 1986).

Most stems collected for the descriptive study (24 for C. haemorrhoidalis and 23 for P. pennipes) contained Odonata eggs. To estimate parasitism rate, we checked 1587 eggs of C. haemorrhoidalis (mean 69.1 \pm 16.8 SE per stem) and 387 eggs of P. pennipes (mean 43 ± 13.7 SE eggs per stem). Parasitoids emerged from 12 stems, with a mean parasitism of $2\% \pm 0.01$ SE for C. haemorrhoidalis and $6\% \pm 0.03$ SE for P. pennipes, and a maximum of 14% and 50%, respectively. In the second study, 551 eggs of C. haemorrhoidalis (mean 26.24 ± 6.7 SE eggs per stem) and 207 eggs of P. pen*nipes* (mean 34.5 ± 12.4 SE eggs per stem) were checked. Parasitoids emerged from 11 stems of a total of 30, with an average parasitism of $8\% \pm 0.03$ SE for C. haemorrhoidalis and $3\% \pm 0.02$ SE for P. pennipes (maximums of 50% and 29%, respectively). There was no significant difference between the overall parasitism rate suffered by both odonates (Mann-Whitney U = 1136.5, P = 0.919).

All parasitoids belonged to the family Mymaridae, and were identified as Anagrus (Anagrus) obscurus Förster 1861, sensu Soyka, 1955 (for anatomical description see Förster 1861; Chiappini 1989, 1996; Baquero & Jordana 1999), a solitary, idiobiont egg parasitoid (Fig. 2). The Mymaridae is a moderately large cosmopolitan family containing nearly 1300 species in about 95 genera. All species are minute (wing length varying from 0.35 to 1.8 mm) solitary (rarely gregarious) endoparasitoids of eggs of Hemiptera, Coleoptera, Odonata and Psocoptera (Gauld & Bolton 1996). They usually parasitize eggs in concealed situations such as those embedded in plant tissues, placed under scales, bracts or in soil (Huber 1986). Adults of some mymarid species spend most of their life under water, being small enough to respire through the general body surface (Askew 1971).

The genus *Anagrus* Haliday is very common in the Palearctic region and well known due to its use in biological control programs (Triapitsyn & Berezovskiy 2004). According to Chiappini (1989), there are 18



Figure 2 Adults of Anagrus obscurus. (A) Female; (B) male.

species of Anagrus in Europe, and all of them are egg parasitoids of Cercopidae, Cicadellidae, Delphacidae, Miridae, Tingidae and Odonata (Huber 1986). Anagrus obscurus was known from Italy as an egg parasitoid of leaf hoppers such as Cicadella viridis L. (Homoptera: Cicadellidae), which lay eggs along the venation of Carex sp. (Cyperaceae) leaves (Chiappini 1989). As far as we know, this is the first time that A. obscurus is reported to be a parasitoid of odonate eggs, and that the parasitoid of C. haemorrhoidalis and P. pennipes is described. A micro-hymenopterous parasitoid of P. pennipes has previously been reported, although the insect remains unidentified (Robert 1958; Martens 1996). In the genus Calopteryx, parasitism of Calopteryx spp. clutches by Aprostocetus (Tetrastichus) natans Kostyukov & Fursov (Hymenoptera, Eulophidae) (Girault 1914), of C. virgo L. by Anagrus incarnatus Haliday (Hymenoptera, Mymaridae) (Henriksen 1919), and of C. atrata Selys by Anaphes sp. (Hymenoptera, Mymaridae) (Asahina 1948) have been reported.

In the laboratory, emerged parasitoids were observed to submerge and walk on Potamogeton stems, apparently searching for host eggs. Some egg parasitoids such as Polynema (Mymaridae) and Tiphodytes (Scelionidae) are known to swim beneath the water to find their hemipteran hosts (Hirashima et al. 1999). The behavior of chalcidoids that locate Odonata hosts above and under water was described by Jarry (1960), who reported that females of A. incarnatus moved over the woody plants where its host, Lestes viridis Vander Linden, had oviposited, repeatedly tapping the bark with the clubbed tips of the antennae to detect host egg presence. Anagrus oviposition occurs only if the host egg is surrounded by living and turgid plant tissues and if dorsal closure of the host embryo is not complete (Witsack 1973).

Parasitism rates suffered by host eggs of both odonates exposed ad libitum (first study) to parasitoid attack or during five days (second study) are quite similar, although the overall number of eggs of C. haemorrhoidalis is almost three times higher in the first study as a consequence of the longer exposition of the oviposition plants. This fact suggests that five days may represent the optimal window of vulnerability of host eggs for A. obscurus. The mechanism by which A. obscurus detects host age is actually unknown, but it is probably related to antennal drumming and sting probing, as other mymarids do (Godfray 1994). Further investigations on host discrimination behavior and host age preference of A. obscurus are needed in the future. Age dependence in the association between parasitoids and hosts, including age dependence in the susceptibility of host to attack, is of potentially great importance in the dynamics of host-parasitoid populations, because host eggs that escape parasitism may represent a demographic refuge that promote stability of the host-parasitoid association (Sait et al. 1995). Although the mean parasitism rate detected is not high, our data suggest that egg parasitoids might be a significant selective factor in the studied locality, and might affect oviposition substrate selection and behavior of egg-laying females of odonates, as has been shown for the water strider Aquarius paludum insularis (Heteroptera: Gerridae), which avoids a high rate of egg parasitism through submerged oviposition on deeper substrates (Amano et al. 2008).

ACKNOWLEDGMENTS

Funding was provided by a grant from the Spanish Ministry of Science and Innovation (CGL2005-00122).

REFERENCES

- Amano H, Hayashi K, Kasuya E (2008) Avoidance of egg parasitism through submerged oviposition by tandem pairs in the water strider, *Aquarius paludum insularis* (Heteroptera: Gerridae). *Ecological Entomology* 33, 560– 563.
- Asahina S (1948) Notes on the odonata of North China. Proceedings of Biogeographical Species of Japan 1, 26–28, 44.
- Askew RR (1971) *Parasitic Insects*. Heinemann Educational Books, London.
- Baquero E, Jordana R (1999) Species of Anagrus Haliday, 1833 (Hymenoptera, Chalcidoidea, Mymaridae) in Navarra (Spain). Miscelània Zoológica 22, 39–50.
- Beckage NE (1985) Endocrine interactions between endoparasitic insects and their hosts. Annual Review of Entomology 30, 371–413.
- Chiappini E (1989) Review of the European species of the genus Anagrus Haliday (Hymenoptera, Chalcidoidea). Bollettino di Zoologia agraria e Bachicoltura 21, 85–119.
- Chiappini E, Triapitsyn SV, Donev A (1996) Key to the Holarctic species of *Anagrus* Haliday (Hymenoptera: Mymaridae) with a review of the Nearctic and Palaearctic (other than European) species and descriptions of new taxa. *Journal of Natural History* 30, 551–595.
- Corbet PS (1999) Dragonflies. Behaviour and Ecology of Odonata. Harley Books, Essex.
- Förster A (1861) Ein Tag in den Hoch-Alpen, eine entomologische Glizze. Programm der Realschule zu Aachen für 1860–1861. 1–44.
- Gauld I, Bolton B (1996) The Hymenoptera. Oxford University Press, Oxford.
- Girault AA (1914) Hosts of insect egg parasitoid in Europe, Asia, Africa and Australasia, with a supplementary American list. Zeitschrift für wissenschaftliche Insektenbiologie 10, 87–91.
- Godfray HCJ (1994) *Parasitoids. Behavioral and Evolutionary Ecology.* Princeton University Press, Princeton, NJ.

- Gross P (1993) Insect behavioral and morphological defenses against parasitoids. *Annual Review of Entomology* 38, 251–273.
- Henriksen KL (1919) De europaeische Vandsnyltehvepse og deres Biologi. Entomologiske Meddelelser 12, 137–251.
- Hirashima Y, Inokuchi M, Yamagishi K (1999) Do you believe a "swimming wasp"? *Esakia* **39**, 9–11.
- Huber JT (1986) Systematics, biology, and hosts of the Mymaridae and Mymarommatidae (Insecta: Hymenoptera). Entomography 4, 185–243.
- Jarry D (1960) Note sur Anagrus incarnatus Hal. (Hymén., Chalcidoïde), parasite des pontes de Lestes viridis L. (Odonate) au Jardin des Plantes. Annales de la Société de Horticulture et Histoire Naturelle d'Herault 100, 59–63.
- Martens A (1996) Die Federlibellen Europas (Platycnemididae). Westarp Wissenschaften, Magdeburg & Spektrum, Heidelberg.
- Robert P-A (1958) Les libellules (odonates). Delachaux et Nestlé, Neuchâtel.
- Sait SM, Andreev A, Begon M, Thompson DJ, Harvey JA, Swain RD (1995) Venturia canescens parasitizing Plodia interpunctella: host vulnerability- a matter of degree. Ecological Entomology 20, 199–201.
- Soyka W (1955) Ueberblick das genus Anagrus Haliday (Alaptidae, Mymaridae, Chalcidoidea, Hymenoptera). Nachrichtenblatt der Bayerischen Entomologen 7, 23–26.
- Strand MR (1986) The physiological interactions of parasitoids with their hosts and their influence on reproductive strategies. In: Waage J, Greathead D (eds) *Insect Parasitoids*, pp 97–136. Academic Press, London.
- Triapitsyn SV, Berezovskiy VV (2004) Review of the genus Anagrus Haliday, 1833 (Hymenoptera: Mymaridae) in Russia, with notes of some extralimital species. Far Eastern Entomologist 139, 1–36.
- Witsack W (1973) Zur Biologie und Ökologie in Zikadeneiern parasitierender Mymariden der Gattung Anagrus (Chalcidoidea, Hymenoptera). Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere 100, 223–299.